



Cihan University/ Sulaymaniyah

College of Health Science

Medical Laboratory Analysis

4th Stage- 1st Semester

Clinical Immunology


Lecture- 1: An Overview of the Immune System (Part-1)

2023- 2024

Lecturer: Mohammed T. Salih


Some Terminology

- **Immune system:** cells, tissues, and molecules that mediate resistance to infections.
- **Immunology:** study of structure and function of the immune system.
- **Immunity:** resistance of a host to pathogens and their toxic effects.
- **Immune response:** collective and coordinated response to the introduction of foreign substances in an individual mediated by the cells and molecules of the immune system.



Immune defense: anti-infection


The Immune System




Functions of the immune system

1. Defense against the growth of tumor cells kills the growth of tumor cells.
2. Protect humans from pathogenic microorganisms.
3. Control of tissue regeneration and scarring.
4. The immune system can injure cells and induce pathologic inflammation.
5. The immune system recognizes and responds to tissue grafts and newly introduced proteins.


Components of the Immune System



Cells of the immune system



Cells and Molecules of the Immune System




Organs of the Immune System



Thymus




Liver



Lymph nodes



Appendix



Tonsil



Red bone marrow




Spleen



Peyer's patches

Types of the Immune System



Factors influencing the level of Innate Immunity

1. Age

- Very old or very young more susceptible to infectious disease.
- Hepatitis B virus infection is asymptomatic in newborn as it lacks developed immune mechanism to express the disease.

2. Hormones

- Endocrine disorders such as **Diabetes Mellitus, hypothyroidism and adrenal dysfunctions** – enhanced susceptibility to infection
- **Increased corticosteroid** – decreased anti-inflammatory effect.

3. Nutrition

- Immune response is reduced in malnutrition patient.




Non- Specific Immunity


First Line of Defense: External Barrier

- General characteristics:
- Most primitive type of immune system found in virtually all multicellular animals.
- High discrimination of host and pathogen
- First line of defense against infection
- No need for prolonged induction.
- Act quickly:
 - Immediate direct response 0-4 hrs.
 - Rapid induced 4-96 hrs.
- Antigen-independent.


Time of the Immune response



Components of Innate Immunity




Barriers to Invasion







Anatomical / Physical/ Mechanical Barriers


System or Organ	Cell type	Mechanism
Skin	Squamous epithelium	Physical barrier (intact skin) Desquamation
Mucous Membranes	Non-ciliated epithelium (<i>e.g.</i> GI tract)	Peristalsis
	Ciliated epithelium, hairs (<i>e.g.</i> respiratory tract)	Mucociliary elevator, Coughing, sneezing
	Epithelium (<i>e.g.</i> nasopharynx)	Flushing action of tears, saliva, mucus, urine; blinking of eye lids

Skin Barrier Function



Normal skin barrier function

- viruses 
- bacteria 
- toxins 
- moisture 




Disrupted skin barrier function


Mucus membrane

Characteristics:

- Membrane that lined the tracts and cavities on the bodies including respiratory, digestive, urinary, reproductive.
- Epithelial cells that secrete mucus, which covers the surface of the mucus membranes.
- Cilia- on mucous membranes- to propel microorganisms out of the body.



Digestive system



Urinary system



Respiratory system

Biological Factors

System or Organ	Component	Mechanism
Skin and mucous membranes	Normal microflora	Antimicrobial substances Competition for nutrients and colonization

Chemical Factors

(At surfaces and in body cavities)

System or Organ	Component	Mechanism
Skin	Sweat	Anti-microbial fatty acids, high salt conc.
Mucous Membranes	HCl (parietal cells) Tears and saliva	Low pH Lysozyme and phospholipase A
	Defensins (respiratory & GI tract)	Antimicrobial
	Sufactants (lung)	Opsonin

Chemical Factors (Humoral Components)

Component	Mechanism
Complement	Lysis of bacteria and some viruses, Opsonin Increase in vascular permeability Recruitment and activation of phagocytic cells
Coagulation system	Increase vascular permeability, Recruitment of phagocytic cells, β -lysin from platelets – a cationic detergent, antibacterial
Acute phase protein	Antibacterial
Lysozyme	Breaks down bacterial cell walls
Lactoferrin and transferrin	Compete with bacteria for iron
Cytokines	Various effects
Interferon	Anti-viral protein


Cellular Components

Cell	Functions
Neutrophils	Phagocytosis and intracellular killing Inflammation and tissue damage
Macrophages	Phagocytosis and intracellular killing Extracellular killing of infected or altered self targets Tissue repair
NK cells	Killing of virus-infected and altered self targets
Eosinophils	Killing of certain parasites


Second Line of Defense:

Internal Defense Line


Inflammation, Phagocytosis, Complement System and Natural Killer cells.



Innate Immune System: Components of Blood




White blood cells (WBCs)



Phagocytes: Neutrophils and Monocytes/ Macrophages

- Immune cells that perform phagocytosis.
- There are two types of circulating phagocytes, neutrophils and monocytes
- These are blood cells that are recruited to sites of infection, where they recognize and ingest microbes for intracellular killing.
- Phagocytes catch the invaders using its cytoplasmic extensions.




Neutrophils

- Neutrophils, also called polymorphonuclear leukocytes (PMNs),
- The most abundant leukocytes in the blood, numbering 4,000 to 10,000 per μL .
- Large numbers released during infection, particularly bacterial and fungal infections,
- Attracted to chemicals released by bacteria also necrotic (dying) tissue during infection and are the first to arrive during an inflammatory response.
- The production of neutrophils is stimulated by cytokines, known as colony-stimulating factors (CSFs),




Neutrophil



Neutrophils



- Unfortunately, this results in the death of the neutrophil as well. as they die after digesting bacteria. Because of this, neutrophils have a lifespan that ranges from a few hours to a few days.
- Dead neutrophils make up a large proportion of pus.
- Also secrete pro-inflammatory cytokines when exposed to inflammatory mediators.
- Every day, 10 billion neutrophils are produced in the bone marrow making them the most abundant white blood cells.




Monocytes

- Less abundant in the blood than neutrophils, numbering 500 to 1000 per μL .
- They also ingest microbes in the blood and in tissues.
- During inflammatory reactions, monocytes enter extravascular tissues and differentiate into cells called macrophages (fixed types),
- Some resident macrophages such as in the brain, liver, and lungs, are derived from progenitors in the yolk sac or fetal liver early during the development of the organism.



MONOCYTE





Macrophages

- From the Greek word meaning ‘big eaters’.
- Macrophages digest the microbes much slower than neutrophils and do not self- destruct during the digestion process.
- Also secrete pro-inflammatory cytokines when exposed to inflammatory mediators.
- Macrophages are more durable and long-lived than neutrophils.
- Life span: months to years
- Most tissues in the body contain resident populations of macrophages that protect the tissue from infection.
- Initiate adaptive immune responses as they display antigens from the pathogens to the lymphocytes (as APC).



SEM of a macrophage white blood cell engulfing bacteria.

In adult homeostasis and Inflammatory reactions

Bone marrow




Hematopoietic stem cell

Monocyte/
dendritic cell
precursor

Blood

Monocyte



During early development

Fetal hematopoietic
organs (yolk sac, liver)



Hematopoietic stem cell

Bone marrow



Hematopoietic stem cell

Tissue
macrophage
precursor

Blood



Differentiation

Tissue

Brain:
Microglial
cells


Liver:
Kupffer cells


Lung:
Alveolar
macrophage

Spleen:
Sinusoidal macrophages

Antigen-presenting cells (APC)


- Antigen-presenting cells (APCs) are a heterogeneous group of immune cells that mediate the cellular immune response by processing and presenting antigens for recognition by T cells.
- Classical APCs include **dendritic cells**, **macrophages**, and **B cells**.
- An antigen-presenting cell (APC) displays foreign antigen complex (parts of the digested invaders) with MHC (Major histocompatibility complex) II on its surface.
- T-cells recognize this complex using their T-cell receptor






Dendritic cells

- More efficient APCs than macrophages.
- These cells reside in the structural compartment of the lymphoid organs such as the thymus, lymph nodes and spleen.
- It can also be found in the bloodstream and other tissues of the body.
- It captures antigen then bring it to the lymphoid organs to start-up the adaptive immune response.




Eosinophils

- These cells have the purpose of fighting parasites such as helminths and in allergic reactions.
- Primary location: intravascular circulation
- Life span: days
- Release pro-inflammatory mediators and toxic-substances against non-phagocytosable surfaces (too large for phagocytes).
- May balance immediate hypersensitivity reactions by degrading or inactivating mediators released by mast cells.



Eosinophils (Cont.)

- The eosinophils release various substances from their eosinophilic granules.
- The granule contents are capable of damaging the parasite membrane.
- It contains many granules rich in:
 1. Major basic protein (MBP)
 2. Cationic proteins
 3. Peroxidase
 4. Arylsulphatase B, phospholipase D
 5. Histaminase
 6. Cytokines




Basophils


- Play a role in host defense, particularly against parasites, and allergic responses.
- Release inflammatory mediators.
- Primary location: intravascular circulation.
- Life span: days
- Contains many granules rich in:
 - 1.Heparin
 - 1.Histamine
 - 2.Serotonin
 - 3.Interleukins

Mast cells

- Play a key role in the inflammatory process and wound healing.
- Its play an important role in immediate hypersensitivity - allergy and anaphylaxis.
- Young basophils settle into tissues and become mast cells.
- Primary location: tissues
- Life span: months
- Contains many granules rich in:
 - 1.Heparin
 - 1.Histamine
 - 2.Serotonin
 - 3.Serine proteases
 - 4.Prostaglandin D2



TEM of a section through a mast cell, with vacuoles (dark) containing histamine or heparin




Natural Killer Cells (NK cells)

- Guard the body against own body cells that are viruses infected or have become cancerous.
- Roaming all over the body looking for abnormal cells to kill.
- NK cells know if our cells became abnormal by scanning their MHC I, abnormal cells that have a different MHC I, or not expressing MHC I, will be killed.
- Life span: weeks
- NK cells will poke and made many holes on the abnormal cells until it died from apoptosis.
- Use cytolytic proteins such as perforin.
- Called natural killer cells because even though they are a type of T cells, they exhibit spontaneous killing of abnormal cells without the need for specific antigen activation as required by T-cells in the adaptive immune response.
- NK cells are important in defense against viral infections and malignancies.

Natural Killer Cells (NK cells)- Cont.

- ❖ NK cells recognize **infected** and **stressed cells** and respond by killing these cells by secreting the **macrophage-activating cytokine IFN- γ** .
- ❖ Activated NK cells empty the contents of their cytoplasmic granules into the extracellular space at the point of contact with the infected cell.
- ❖ The granule proteins then enter infected cells and activate enzymes that induce **apoptosis**.



Second Line of Defense: Internal Defense Line

Inflammatory Response, Complement System




Inflammatory response

- Activate the body 'alarm' using chemicals.
- Biological response of the immune system that can be triggered by a variety of factors:
 1. Pathogens,
 2. Damaged cells,
 3. Toxic compounds,



Inflammation




Inflammation has two main components

1. The cellular component involves the movement of white blood cells from blood vessels into the inflamed tissue. extravasate (filter out) phagocytes, picking up bacteria and cellular debris.
2. The exudative component involves the movement of fluid, usually containing many important proteins such as fibrin and immunoglobulins (antibodies).

The Signs and Symptoms Include:

- Redness (Rubor)- raised blood flow.
- Swelling (Tumor)– leakage of fluids to the tissue.
- Heat (Calor) – raised blood flow, leakage of fluids to the tissue, release of inflammatory mediators.
- Pain (Dolor)– stimulation of pain receptors by inflammatory mediators, injury of nerve fibres, and irritation by toxic chemicals released by microbes.
- To some extent, loss of tissue function (Functio laesa)– pain, disruption of tissue structure, fibroplasia, metaplasia.


5 Cardinal Signs of Inflammation



Pain



Heat



Redness




Swelling



**Loss of
Function**


1- Release of soluble mediators



- An injury site exposed to invaders.
- Mast cells and the injured vascular epithelium will set up the inflammatory response by releasing histamine and other vasoactive substances.

The tissue damage triggers mast cells to release histamine from their histamine granules into the surrounding tissue.


2- Vasodilation



- Histamine main effect is to activate vasodilation, which leads to the leakage of vascular fluid to the site of injury.

Histamine diffuses into capillaries, causing them to dilate and become leaky. As capillaries dilate, the area appears red. As plasma leaks into the tissue, the area swells.


3- Increased blood flow and Extravasation



In damaged or infected tissue, activated proteins called complement proteins and other chemicals attract phagocytes to the area.

1. The reason is to flood the area with serum proteins that contain:
2. Inflammatory mediators (prostaglandins, kininogen, complement, cytokines etc.)
3. Antimicrobial proteins (defensins, hepcidins, cathelicidin etc.)
4. Chemoattractant (C3a, C5a, leukotriene etc.)

4- Migration of Phagocytes (Chemotaxis)




The phagocytes engulf and digest dead cells and bacteria.

Monocytes, neutrophils and other effector cells from the blood attracted to the sites of infection or tissue damage by chemo-attractants.

Helps by the:

- Increased margination (the gap between the cells)
- Leakage of serum fluids to the infected area.

5- Tissue Repair



Inflammatory response continues until the foreign material is eliminated and the injury is repaired.

Histamine and complement signaling cease, and phagocytes are no longer attracted to the area. The tissue returns to normal.


Benefits of the inflammatory response

- Protective Role:
 1. Prevents the spread of damaging agents to nearby tissues,
 2. Clearance of foreign or damaged material,
 3. Encourages repair,
- The downside of the inflammatory reaction is that it is nonspecific and the inflammatory compounds released are also harmful to the host.
- Because of this, inflammation is under tight regulatory control in the body.

Phagocytosis

- Phago = to eat
- Cyte = cell
- WBCs (eg. Neutrophils) – find, eat and digest microbes!
- The process by which phagocytes ingest or engulf other cells or particles.
- Resting phagocytes become activated by inflammatory mediators.
- This activated full of energy phagocytes moves towards the chemical attractant (chemoattractant).
- Majority of the eaten microbes will be digested into debris, while small parts of it will be processed then presented into adaptive immune cells.

Phases of Phagocytosis



- 1 Chemotaxis and adherence of microbe to phagocyte.
- 2 Ingestion of microbe by phagocyte.
- 3 Formation of a phagosome.
- 4 Fusion of the phagosome with a lysosome to form a phagolysosome.
- 5 Digestion of ingested microbe by enzymes.
- 6 Formation of residual body containing indigestible material.
- 7 Discharge of waste materials.

Granules

What's in the granules ? Lysozyme – digests bacterial cell wall; other antimicrobial proteins.


Serum proteins: Complement (C')

- They are a large number of distinct plasma proteins produced by the liver that react with one another (C1 thro' C9)
- Circulated in blood plasma and tissues
- System of plasma proteins that can be activated:
 - ✓ Directly by pathogens
 - ✓ Indirectly by pathogen-bound antibody
- Activation leading to a cascade of reactions that occurs on the surface of pathogens and generates active components with various effector functions
- Complement can bind to microbes and coat the microbes,
- Essential part of innate immune response,
- Enhances adaptive immune response.


The Functions of Complement

1. Lysis of cells, bacteria, and viruses,
2. Opsonization, which promotes phagocytosis of particulate antigens,
3. Binding to specific complement receptors on cells of the immune system, triggering specific cell functions, inflammation, and secretion of immunoregulatory molecules,
4. Immune clearance, which removes immune complexes from the circulation and deposits them in the spleen and liver.


Complement Proteins



Complement proteins: Role in innate immune system






How do Complement Proteins Facilitate Phagocytosis ?



How do Complement Proteins Lyse Pathogens?

Membrane attack complex formed by c` proteins



① Activated complement proteins form complexes of proteins that create holes in the bacterial cell wall.


② Water and salts diffuse into the bacterium through the holes.

③ The bacterium swells and eventually bursts.

Copyright © 2001 Benjamin Cummings, an imprint of Addison Wesley Longman, Inc.

References

1. Turgeon, M. L. (1996). Immunology & serology in laboratory medicine. St. Louis: Mosby.
2. Abul K. Abbas and Andrew H. Lichtman. Cellular And Molecular Immunology 2019, 6th edition .
3. **Abbas, A. K., Lichtman, A. H., Pillai, S., & Baker, D. L. (. i. (2020).** Basic immunology: Functions and disorders of the immune system (Sixth edition.).
4. Aryal (2021) Cells Of The Immune System. <https://microbenotes.com/cells-of-the-immune-system/>
5. British society for Immunology_ What is immunology?
<https://www.immunology.org/public-information/what-is-immunology#:~:text=Immunology%20is%20the%20study%20of,as%20autoimmunity%2C%20allergy%20and%20cancer>



Here is the required reading for this lecture