University of Cihan-Sulaimaniya College of Science Department of MLA Second year student



Human Anatomy

Week Eight

Cardiovascular System

1 - 11 -2023

- The **heart** is a conical hollow muscular organ situated in the middle mediastinum and is enclosed within the pericardium. It is positioned posteriorly to the body of the sternum with one-third situated on the right and two-thirds on the left of the midline.
- The heart is laid on its right side



Pericardium

Is the outermost layer of fibroserous sac that encloses the heart and the roots of the great vessels



serosa.

Gross anatomy of the heart



Original Impulses from S-A Node

The electrical impulses are normally generated by a group of specialized pacemaker cells at *sinoatrial (SA) node*.



- Applications of ECG
- 1) Measure automaticity HR, rhythmicity, pacemaker
- 2) Measure conductivity pathway, reentry, block
- 3) Reveal hypertrophy
- 4) Reveal ischemic damages location, size, and progress

Bradycardia Slow heart rate (< 60 beats/min) *Tachycardia* Fast heart rate (> 100 beats/min)



Major blood vessels of the heart



The cardiac muscles get nutrients from coronary circulation.

Coronary Vessels — Posterior Aspect



Coronary Vessels — Anterior Aspect





- Stimulated by increase in arterial pressure (stretch)
- Regulate the heart when blood pressure (BP) increases or drops
- Involved in short term regulation of BP

Chemoreceptor Reflex



systemic circulation

At the arch of the aorta, 3 branches extend upward;

- The <u>brachiocephalic artery</u>, quickly divides into the <u>right subclavian artery</u> that supplies blood to the right arm and upper torso and the <u>right common</u> <u>carotid artery</u> that supplies the head and neck.
- 2. The <u>left common carotid artery</u> supplies the head and neck.
- 3. The <u>left subclavian artery</u> supplies the left arm and upper torso. 'Subclavian' means it is located below the clavicle... or collarbone









Main Veins







Copyright © 2001 Benjamin Cummings, an imprint of Addison Wesley Longman, Inc.

LYMPHATIC SYSTEM

Between the cells of the body are spaces filled with fluid. This is the interstitial (or tissue) fluid, often referred to as intercellular fluid.

There are continuous exchanges between the intracellular fluid, the interstitial fluid, and the plasma of the blood.

The lymphatic system returns to the bloodstream the excess interstitial fluid, which includes proteins and fluid derived from the blood.

- a) Lymphatic Capillaries. Lymphatic capillaries are located in the interstitial spaces. Here, they absorb the excess fluids.
- b) b. Lymph Vessels. A tributary system of vessels collects these excess fluids (lymph)
- c) Lymph Nodes. Along the way, lymphatic vessels are interrupted by special structures known as lymph nodes.
- d) Tonsils. Tonsils are special collections of lymphoid tissue, very similar to a group of lymph nodes.

